

# MATHEMATICS

## Chapter 6: Triangles




## Triangles

### 1. Congruent figures:


Two geometrical figures are called **congruent** if they superpose exactly on each other, that is, they are of the same shape and size.

### 2. Similar figures:


Two figures are **similar**, if they are of the same shape but not necessarily of the same size.



Group 1



Group 2



Group 3

3. All congruent figures are similar but the similar figures need not to be congruent.

4. Two **polygons** having the same number of sides are **similar** if

- i. their corresponding angles are equal and
- ii. their corresponding sides are in the same ratio (or proportion).

**Note:** Same ratio of the corresponding sides means the **scale factor** for the polygons.

5. Important facts related to similar figures are:

- i. All circles are similar.
- ii. All squares are similar.
- iii. All equilateral triangles are similar.
- iv. The ratio of any two corresponding sides in two equiangular triangles is always same.

6. Two **triangles are similar** ( $\sim$ ) if

- i. Their corresponding angles are equal.
- ii. Their corresponding sides are in same ratio.


7. If the angles in two triangles are:

- i. Different, the triangles are neither similar nor congruent.
- ii. Same, the triangles are similar.
- iii. Same and the corresponding sides are of the same size, the triangles are congruent.

In the given figure,  $A \leftrightarrow D$ ,  $B \leftrightarrow E$  and  $C \leftrightarrow F$ , which means triangles ABC and DEF are similar which is represented by  $\Delta ABC \sim \Delta DEF$







8. If  $\Delta ABC \sim \Delta PQR$ , then

- i.  $\angle A = \angle P$
- ii.  $\angle B = \angle Q$
- iii.  $\angle C = \angle R$
- iv.  $\frac{AB}{PQ} = \frac{BC}{QR} = \frac{AC}{PR}$

#### 9. Equiangular triangles:

Two triangles are **equiangular** if their corresponding angles are equal. The ratio of any two corresponding sides in such triangles is always the same.

#### 10. Basic Proportionality Theorem (Thales Theorem):

If a line is drawn parallel to one side of a triangle to intersect other two sides in distinct points, the other two sides are divided in the same ratio.

#### 11. Converse of BPT:

If a line divides any two sides of a triangle in the same ratio then the line is parallel to the third side.

12. A line drawn through the mid-point of one side of a triangle which is parallel to another side bisects the third side. In other words, the line joining the mid-points of any two sides of a triangle is parallel to the third side.

#### 13. AAA (Angle-Angle-Angle) similarity criterion:

If in two triangles, corresponding angles are equal, then their corresponding sides are in the same ratio (or proportion) and hence the two triangles are similar.

#### 14. AA (Angle-Angle) similarity criterion:

If two angles of a triangle are respectively equal to two angles of another triangle, then by the angle sum property of a triangle their third angles will also be equal.

Thus, **AAA similarity criterion** changes to **AA similarity criterion** which can be stated as follows:

If two angles of one triangle are respectively equal to two angles of other triangle, then the two triangles are similar.

#### 15. Converse of AAA similarity criterion:

If two triangles are similar, then their corresponding angles are equal.

#### 16. SSS (Side-Side-Side) similarity criterion:

If in two triangles, sides of one triangle are proportional to (i.e., in the same ratio of) the sides of the other triangle, then their corresponding angles are equal and hence the two



triangles are similar.

### 17. Converse of SSS similarity criterion:

If two triangles are similar, then their corresponding sides are in constant proportion.

### 18. SAS (Side-Angle-Side) similarity criterion:

If one angle of a triangle is equal to one angle of the other triangle and the sides including these angles are proportional, then the two triangles are similar.

### 19. Converse of SAS similarity criterion:

If two triangles are similar, then one of the angles of one triangle is equal to the corresponding angle of the other triangle and the sides including these angles are in constant proportion.

### 20. RHS (Right angle-Hypotenuse-Side) criterion:

If in two right triangles, hypotenuse and one side of one triangle are proportional to the hypotenuse and one side of another triangle, then the two triangles are similar. This criteria is referred as the RHS similarity criterion

### 21. Pythagoras Theorem:

In a right triangle, the square of the hypotenuse is equal to the sum of the squares of the other two sides.

Thus, in triangle  $ABC$  right angled at  $B$ ,  $AB^2 + BC^2 = AC^2$

### 22. Converse of Pythagoras Theorem:

If in a triangle, square of one side is equal to the sum of the squares of the other two sides, then the angle opposite the first side is a right angle.

### 23. The ratio of the areas of two similar triangles is equal to the square of the ratio of their corresponding sides.

Thus, if  $\Delta ABC \sim \Delta PQR$ , then  $\frac{ar \Delta ABC}{ar \Delta PQR} = \left(\frac{AB}{PQ}\right)^2 = \left(\frac{BC}{QR}\right)^2 = \left(\frac{CA}{RP}\right)^2$

Also, the ration of the areas of two similar triangles is equal to the ration of the squares of the corresponding medians.

### 24. Some important results of similarity are:

In an equilateral or an isosceles triangle, the altitude divides the base into two equal parts.

If a perpendicular is drawn from the vertex of the right triangle to the hypotenuse then triangles on both sides of the perpendicular are similar to the whole triangle and to each other.

The area of an equilateral triangle described on one side of a square is equal to half the area of the equilateral triangle described on one of its diagonal.

Sum of the squares of the sides of a rhombus is equal to the sum of the squares of its diagonals.



In an equilateral triangle, three times the square of one side is equal to four times the square of one of its altitudes.

## 25. Triangle

A triangle can be defined as a polygon which has three angles and three sides. The interior angles of a triangle sum up to 180 degrees and the exterior angles sum up to 360 degrees. Depending upon the angle and its length, a triangle can be categorized in the following types-

- Scalene Triangle – All the three sides of the triangle are of different measure
- Isosceles Triangle – Any two sides of the triangle are of equal length
- Equilateral Triangle – All the three sides of a triangle are equal and each angle measures 60 degrees
- Acute angled Triangle – All the angles are smaller than 90 degrees
- Right angle Triangle – Anyone of the three angles is equal to 90 degrees
- Obtuse-angled Triangle – One of the angles is greater than 90 degrees

## 26. Similarity Criteria of Triangles

To find whether the given two triangles are similar or not, it has four criteria. They are:

**Side-Side- Side (SSS) Similarity Criterion** – When the corresponding sides of any two triangles are in the same ratio, then their corresponding angles will be equal and the triangle will be considered as similar triangles.

**Angle Angle Angle (AAA) Similarity Criterion** – When the corresponding angles of any two triangles are equal, then their corresponding side will be in the same ratio and the triangles are considered to be similar.

**Angle-Angle (AA) Similarity Criterion** – When two angles of one triangle are respectively equal to the two angles of the other triangle, then the two triangles are considered as similar.

**Side-Angle-Side (SAS) Similarity Criterion** – When one angle of a triangle is equal to one angle of another triangle and the sides including these angles are in the same ratio (proportional), then the triangles are said to be similar.

## 27. Proof of Pythagoras Theorem

Statement: As per Pythagoras theorem, “In a right-angled triangle, the sum of squares of two sides of a right triangle is equal to the square of the hypotenuse of the triangle.”

Proof –


Consider the right triangle, right-angled at B.

Construction-





Draw  $BD \perp AC$



Now,  $\triangle ADB \sim \triangle ABC$

So,  $AD/AB = AB/AC$

or  $AD \cdot AC = AB^2$  .....(i)

Also,  $\triangle BDC \sim \triangle ABC$

So,  $CD/BC = BC/AC$

or,  $CD \cdot AC = BC^2$  .....(ii)

Adding (i) and (ii),

$AD \cdot AC + CD \cdot AC = AB^2 + BC^2$

$AC (AD + DC) = AB^2 + BC^2$

$AC (AC) = AB^2 + BC^2$

$\Rightarrow AC^2 = AB^2 + BC^2$

Hence, proved.

## 28. Problems Related to Triangles

A girl having a height of 90 cm is walking away from a lamp-post's base at a speed of 1.2 m/s. Calculate the length of that girl's shadow after 4 seconds if the lamp is 3.6 m above the ground.

S and T are points on sides PR and QR of triangle PQR such that  $\angle P = \angle RTS$ . Now, prove that triangle RPQ and triangle RTS are similar.


E is a point on the side AD produced of a parallelogram ABCD and BE intersects CD at F. Show that triangles ABE and CFB are similar.

## 29. Pythagoras Theorem Statement

Pythagoras theorem states that "In a right-angled triangle, the square of the hypotenuse side is equal to the sum of squares of the other two sides". The sides of this triangle have been named as Perpendicular, Base and Hypotenuse. Here, the hypotenuse is the longest side, as it is opposite to the angle  $90^\circ$ . The sides of a right triangle (say a, b and c) which have positive integer values, when squared, are put into an equation, also called a Pythagorean triple.







### History

The theorem is named after a greek Mathematician called Pythagoras.

Pythagoras Theorem Formula

Consider the triangle given above:

Where “a” is the perpendicular,

“b” is the base,


“c” is the hypotenuse.

According to the definition, the Pythagoras Theorem formula is given as:

$$\text{Hypotenuse}^2 = \text{Perpendicular}^2 + \text{Base}^2$$

$$c^2 = a^2 + b^2$$

The side opposite to the right angle (90°) is the longest side (known as Hypotenuse) because the side opposite to the greatest angle is the longest.



Consider three squares of sides a, b, c mounted on the three sides of a triangle having the same sides as shown.



By Pythagoras Theorem –


Area of square “a” + Area of square “b” = Area of square “c”

### Pythagoras Theorem Proof

Given: A right-angled triangle ABC, right-angled at B.

To Prove-  $AC^2 = AB^2 + BC^2$

Construction: Draw a perpendicular BD meeting AC at D.



Proof:

We know,  $\triangle ADB \sim \triangle ABC$

Therefore,

$$\frac{AD}{AB} = \frac{AB}{AC}$$

(corresponding sides of similar triangles)

$$\text{Or, } AB^2 = AD \times AC \dots\dots\dots(1)$$

Also,  $\triangle BDC \sim \triangle ABC$

Therefore,

$$\frac{CD}{BC} = \frac{BC}{AC}$$

(corresponding sides of similar triangles)

$$\text{Or, } BC^2 = CD \times AC \dots\dots\dots(2)$$

Adding the equations (1) and (2) we get,

$$AB^2 + BC^2 = AD \times AC + CD \times AC$$


$$AB^2 + BC^2 = AC (AD + CD)$$

Since,  $AD + CD = AC$

$$\text{Therefore, } AC^2 = AB^2 + BC^2$$

Hence, the Pythagorean theorem is proved.





## Important Questions

### Multiple Choice questions-

1. If in triangles ABC and DEF,  $\frac{AB}{EF} = \frac{AC}{DE}$ , then they will be similar when

- (a)  $\angle A = \angle D$
- (b)  $\angle A = \angle E$
- (c)  $\angle B = \angle E$
- (d)  $\angle C = \angle F$

2. A square and a rhombus are always

- (a) similar
- (b) congruent
- (c) similar but not congruent
- (d) neither similar nor congruent

3. If  $\triangle ABC \sim \triangle DEF$  and  $EF = \frac{1}{3} BC$ , then  $ar(\triangle ABC) : ar(\triangle DEF)$  is

- (a) 3 : 1.
- (b) 1 : 3.
- (c) 1 : 9.
- (d) 9 : 1.

4. If a triangle and a parallelogram are on the same base and between same parallels, then what is the ratio of the area of the triangle to the area of parallelogram?

- (a) 1 : 2
- (b) 3 : 2
- (c) 1 : 3
- (d) 4 : 1

5. D and E are respectively the points on the sides AB and AC of a triangle ABC such that  $AD = 2$  cm,  $BD = 3$  cm,  $BC = 7.5$  cm and  $DE \parallel BC$ . Then, length of DE (in cm) is





(a) 2.5

(b) 3

(c) 5

(d) 6

6. Which geometric figures are always similar?

(a) Circles

(b) Circles and all regular polygons

(c) Circles and triangles

(d) Regular

7.  $\triangle ABC \sim \triangle PQR$ ,  $\angle B = 50^\circ$  and  $\angle C = 70^\circ$  then  $\angle P$  is equal to

(a)  $50^\circ$

(b)  $60^\circ$

(c)  $40^\circ$

(d)  $70^\circ$

8. In triangle DEF, GH is a line parallel to EF cutting DE in G and DF in H. If DE = 16.5, DH = 5, HF = 6 then GE = ?

(a) 9

(b) 10

(c) 7.5

(d) 8

9. In a rectangle Length = 8 cm, Breadth = 6 cm. Then its diagonal = ...

(a) 9 cm

(b) 14 cm

(c) 10 cm

(d) 12 cm



10. In triangle ABC,  $DE \parallel BC$   $AD = 3$  cm,  $DB = 8$  cm  $AC = 22$  cm. At what distance from A does the line DE cut AC?

- (a) 6
- (b) 4
- (c) 10
- (d) 5

### Very Short Questions:

1. Two sides and the perimeter of one triangle are respectively three times the corresponding sides and the perimeter of the other triangle. Are the two triangles similar? Why?
2. A and B are respectively the points on the sides PQ and PR of a  $\Delta PQR$  such that  $PQ = 12.5$  cm,  $PA = 5$  cm,  $BR = 6$  cm, and  $PB = 4$  cm. Is  $AB \parallel QR$ ? Give reason.
3. If  $\Delta ABC \sim \Delta QRP$ ,  $\frac{ar(\Delta ABC)}{ar(\Delta PQR)} = \frac{9}{4}$ ,  $AB = 18$  cm and  $BC = 15$  cm, then find the length of PR.
4. If it is given that  $\Delta ABC \sim \Delta PQR$  with  $\frac{BC}{QR} = \frac{1}{3}$ , then find  $\frac{ar(\Delta PQR)}{ar(\Delta ABC)}$
5.  $\Delta DEF \sim \Delta ABC$ , if  $DE : AB = 2 : 3$  and  $ar(\Delta DEF)$  is equal to 44 square units. Find the area ( $\Delta ABC$ ).
6. Is the triangle with sides 12 cm, 16 cm and 18 cm a right triangle? Give reason.
7. In triangles PQR and TSM,  $\angle P = 55^\circ$ ,  $\angle Q = 25^\circ$ ,  $\angle M = 100^\circ$ , and  $\angle S = 25^\circ$ . Is  $\Delta QPR \sim \Delta TSM$ ? Why?
8. If ABC and DEF are similar triangles such that  $\angle A = 47^\circ$  and  $\angle E = 63^\circ$ , then the measures of  $\angle C = 70^\circ$ . Is it true? Give reason.
9. Let  $\Delta ABC \sim \Delta DEF$  and their areas be respectively  $64 \text{ cm}^2$  and  $121 \text{ cm}^2$ . If  $EF = 15.4$  cm, find BC.
10. ABC is an isosceles triangle right-angled at C. Prove that  $AB^2 = 2AC^2$ .

### Short Questions :

1. In Fig. 7.10,  $DE \parallel BC$ . If  $AD = x$ ,  $DB = x - 2$ ,  $AE = x + 2$  and  $EC = x - 1$ , find the value of x.







Fig. 7.10

- E and F are points on the sides PQ and PR respectively of a  $\Delta PQR$ . Show that  $EF \parallel QR$  if  $PQ = 1.28$  cm,  $PR = 2.56$  cm,  $PE = 0.18$  cm and  $PF = 0.36$  cm.
- A vertical pole of length 6 m casts a shadow 4 m long on the ground and at the same time a tower casts a shadow 28 m long. Find the height of the tower.
- In Fig. 7.13, if  $LM \parallel CB$  and  $LN \parallel CD$ , prove that  $\frac{AM}{AB} = \frac{AN}{AD}$
- In Fig. 7.14,  $DE \parallel OQ$  and  $DF \parallel OR$  Show that  $EF \parallel QR$ .
- Using converse of Basic Proportionality Theorem, prove that the line joining the mid-points of any two sides of a triangle is parallel to the third side.
- State which pairs of triangles in the following figures are similar. Write the similarity criterion used by you for answering the question and also write the pairs of similar triangles in the symbolic form.



- In Fig. 7.17,  $\frac{AO}{OC} = \frac{BO}{OD} = \frac{1}{2}$  and  $AB = 5$  cm. Find the value of  $DC$ .
- E is a point on the side AD produced of a parallelogram ABCD and BE intersects CD at F. Show that  $\Delta ABE \sim \Delta CFB$ .
- S and T are points on sides PR and QR of  $\Delta PQR$  such that  $\angle P = \angle RTS$ . Show that  $\Delta RPQ \sim \Delta RTS$ .

### Long Questions :

- Using Basic Proportionality Theorem, prove that a line drawn through the mid-point of one side of a triangle parallel to another side bisects the third side.
- ABCD is a trapezium in which  $AB \parallel DC$  and its diagonals intersect each other



at the point O. Show that  $\frac{AO}{BO} = \frac{CO}{DO}$ .

3. If AD and PM are medians of triangles ABC and PQR respectively, where  $\Delta ABC \sim \Delta PQR$ , prove that  $\frac{AB}{PQ} = \frac{AD}{PM}$
4. In Fig. 7.37, ABCD is a trapezium with  $AB \parallel DC$ . If  $\Delta AED$  is similar to  $\Delta BEC$ , prove that  $AD = BC$ .
5. Prove that the area of an equilateral triangle described on a side of a right-angled isosceles triangle is half the area of the equilateral triangle described on its hypotenuse.
6. If the areas of two similar triangles are equal, prove that they are congruent.
7. Prove that the ratio of the areas of two similar triangles is equal to the square of the ratio of their corresponding medians.
8. In Fig. 7.41, O is a point in the interior of a triangle ABC,  $OD \perp BC$ ,  $OE \perp AC$  and  $OF \perp AB$ . Show that
  - (i)  $OA^2 + OB^2 + OC^2 - OD^2 - OE^2 - OF^2 = AF^2 + BD^2 + CE^2$
  - (ii)  $AF^2 + BD^2 + CE^2 = AE^2 + CD^2 + BF^2$
9. The perpendicular from A on side BC of a  $\Delta ABC$  intersects BC at D such that  $DB = 3CD$  (see Fig. 7.42). Prove that  $2AB^2 = 2AC^2 + BC^2$
10. In an equilateral triangle, prove that three times the square of one side is equal to four times the square of one of its altitudes.

### Case Study Questions:



1. Rahul is studying in X Standard. He is making a kite to fly it on a Sunday. Few questions came to his mind while making the kite. Give answers to his questions by looking at the




figure.

- i. Rahul tied the sticks at what angles to each other?
  - a.  $30^\circ$
  - b.  $60^\circ$
  - c.  $90^\circ$
  - d.  $60^\circ$
- ii. Which is the correct similarity criteria applicable for smaller triangles at the upper part of this kite?
  - a. RHS
  - b. SAS
  - c. SSA
  - d. AAS
- iii. Sides of two similar triangles are in the ratio 4:9. Corresponding medians of these triangles are in the ratio:
  - a. 2 : 3
  - b. 4 : 9
  - c. 81 : 16
  - d. 16 : 81
- iv. In a triangle, if the square of one side is equal to the sum of the squares of the other two sides, then the angle opposite the first side is a right angle. This theorem is called.
  - a. Pythagoras theorem



- b. Thales theorem
    - c. The converse of Thales theorem
    - d. The converse of Pythagoras theorem
  - v. What is the area of the kite, formed by two perpendicular sticks of length 6cm and 8cm?
    - a.  $48 \text{ cm}^2$
    - b.  $14 \text{ cm}^2$
    - c.  $24 \text{ cm}^2$
    - d.  $96 \text{ cm}^2$
2. There is some fire incident in the house. The fireman is trying to enter the house from the window as the main door is locked. The window is 6m above the ground. He places a ladder against the wall such that its foot is at a distance of 2.5m from the wall and its top reaches the window.



- i. Here, \_\_\_\_\_ be the ladder and \_\_\_\_\_ be the wall with the window.
  - a. CA, AB
  - b. AB, AC
  - c. AC, BC
  - d. AB, BC
- ii. We will apply Pythagoras Theorem to find length of the ladder. It is:
  - a.  $AB^2 = BC^2 - CA^2$
  - b.  $CA^2 = BC^2 + AB^2$
  - c.  $BC^2 = AB^2 + CA^2$

d.  $AB^2 = BC^2 + CA^2$

- iii. The length of the ladder is \_\_\_\_\_.
- a. 4.5m
  - b. 2.5m
  - c. 6.5m
  - d. 5.5m
- iv. What would be the length of the ladder if it is placed 6m away from the wall and the window is 8m above the ground?
- a. 12m
  - b. 10m
  - c. 14m
  - d. 8m
- v. How far should the ladder be placed if the fireman gets a 9m long ladder?
- a. 6.7m (approx.)
  - b. 7.7m (approx.)
  - c. 5.7m (approx.)
  - d. 4.7m (approx.)

### Assertion Reason Questions-

1. **Directions:** In the following questions, a statement of assertion (A) is followed by a statement of reason (R). Mark the correct choice as:
- a. Both assertion (A) and reason (R) are true and reason (R) is the correct explanation of assertion (A).
  - b. Both assertion (A) and reason (R) are true but reason (R) is not the correct explanation of assertion (A).
  - c. Assertion (A) is true but reason (R) is false.
  - d. Assertion (A) is false but reason (R) is true.

**Assertion:** If two sides of a right angle are 7 cm and 8 cm, then its third side will be 9 cm.

**Reason:** In a right triangle, the square of the hypotenuse is equal to the sum of the squares of the other two sides.



2. **Directions:** In the following questions, A statement of Assertion (A) is followed by a statement of Reason (R). Mark the correct choice as.
- a. Both assertion (A) and reason (R) are true and reason (R) is the correct explanation of assertion (A).
  - b. Both assertion (A) and reason (R) are true but reason (R) is not the correct explanation of assertion (A).
  - c. Assertion (A) is true but reason (R) is false.
  - d. Assertion (A) is false but reason (R) is true.

**Assertion:** If  $\triangle ABC$  and  $\triangle PQR$  are congruent triangles, then they are also similar triangles.

**Reason:** All congruent triangles are similar but the similar triangles need not be congruent.





Answer Key-**Multiple Choice questions-**

1. (b)  $\angle A = \angle E$
2. (d) neither similar nor congruent
3. (c) 1 : 9.
4. (a) 1 : 2
5. (b) 3
6. (b) Circles and all regular polygons
7. (b)  $60^\circ$
8. (a) 9
9. (c) 10cm
10. (a) 6

**Very Short Answer :**

1. Since the perimeters and two sides are proportional  
 $\therefore$  The third side is proportional to the corresponding third side.  
 i.e., The two triangles will be similar by SSS criterion.

2.

Yes,  $\frac{PA}{AQ} = \frac{5}{12.5 - 5} = \frac{5}{7.5} = \frac{2}{3}$

$$\frac{PB}{BR} = \frac{4}{6} = \frac{2}{3}$$

Since  $\frac{PA}{AQ} = \frac{PB}{BR} = \frac{2}{3}$

$\therefore AB \parallel QR$




Fig. 7.4

3.

$$\frac{\text{Area of } \triangle ABC}{\text{Area of } \triangle QRP} = \frac{BC^2}{RP^2} \Rightarrow \frac{9}{4} = \frac{(15)^2}{RP^2}$$

$$\therefore RP^2 = \frac{225 \times 4}{9} = \frac{900}{9} = 100 \Rightarrow RP = 10 \text{ cm}$$

4.

$$\frac{BC}{QR} = \frac{1}{3} \quad (\text{Given})$$

$$\frac{\text{ar}(\triangle PQR)}{\text{ar}(\triangle ABC)} = \frac{(QR)^2}{(BC)^2} \quad [\because \text{Ratio of area of similar triangles is equal to the ratio of square of its corresponding sides}]$$

$$= \left(\frac{QR}{BC}\right)^2 = \left(\frac{3}{1}\right)^2 = \frac{9}{1} = 9 : 1$$

5.

Since  $\triangle DEF \sim \triangle ABC$ 

$$\frac{\text{ar}(\triangle DEF)}{\text{ar}(\triangle ABC)} = \frac{(DE)^2}{(AB)^2}$$

[ $\because$  Ratio of area of similar triangles is equal to the ratio of square of its corresponding sides]

$$\frac{44}{\text{ar}(\triangle ABC)} = \left(\frac{2}{3}\right)^2 \Rightarrow \text{ar}(\triangle ABC) = \frac{44 \times 9}{4}$$

$$\text{So, ar}(\triangle ABC) = 99 \text{ cm}^2$$

$$6. \text{ Here, } 12^2 + 16^2 = 144 + 256 = 400 \neq 182$$

$\therefore$  The given triangle is not a right triangle.

$$7. \text{ Since, } \angle R = 180^\circ - (\angle P + \angle Q)$$

$$= 180^\circ - (55^\circ + 25^\circ) = 100^\circ = \angle M$$

$$\angle Q = \angle S = 25^\circ \text{ (Given)}$$

$$\triangle QPR \sim \triangle STM$$

i.e.,  $\triangle QPR$  is not similar to  $\triangle TSM$ .

$$8. \text{ Since } \triangle ABC \sim \triangle DEF$$

$$\therefore \angle A = \angle D = 47^\circ$$

$$\angle B = \angle E = 63^\circ$$



$$\therefore \angle C = 180^\circ - (\angle A + \angle B) = 180^\circ - (47^\circ + 63^\circ) = 70^\circ$$

$\therefore$  Given statement is true.

9.

We have,  $\frac{\text{area of } \triangle ABC}{\text{area of } \triangle DEF} = \frac{BC^2}{EF^2} = (\text{as } \triangle ABC \sim \triangle DEF)$

$$\Rightarrow \frac{64}{121} = \frac{BC^2}{EF^2} \Rightarrow \frac{64}{121} = \frac{BC^2}{(15.4)^2}$$

$$\Rightarrow \frac{BC}{15.4} = \frac{8}{11}$$

$$\therefore BC = \frac{8}{11} \times 15.4 = 11.2 \text{ cm}$$




Fig. 7.5

10.  $\triangle ABC$  is right-angled at C.

$$\therefore AB^2 = AC^2 + BC^2 \text{ [By Pythagoras theorem]}$$

$$\Rightarrow AB^2 = AC^2 + AC^2$$

$$[\because AC = BC]$$

$$\Rightarrow AB^2 = 2AC^2$$

### Short Answer :

1. In  $\triangle ABC$ , we have

$$DE \parallel BC,$$

$$\therefore \frac{AD}{DB} = \frac{AE}{EC} \text{ [By Basic Proportionality Theorem]}$$

$$\Rightarrow \frac{x}{x-2} = \frac{x+2}{x-1}$$

$$\Rightarrow x(x-1) = (x-2)(x+2)$$

$$\Rightarrow x^2 - x = x^2 - 4$$

$$\Rightarrow x = 4$$

2.






Fig. 7.11

We have,  $PQ = 1.28$  cm,  $PR = 2.56$  cm

$PE = 0.18$  cm,  $PF = 0.36$  cm

Now,  $EQ = PQ - PE = 1.28 - 0.18 = 1.10$  cm and

$FR = PR - PF = 2.56 - 0.36 = 2.20$  cm

$$\text{Now, } \frac{PE}{EQ} = \frac{0.18}{1.10} = \frac{18}{110} = \frac{9}{55}$$

$$\text{and, } \frac{PF}{FR} = \frac{0.36}{2.20} = \frac{36}{220} = \frac{9}{55} \quad \therefore \frac{PE}{EQ} = \frac{PF}{FR}$$

Therefore,  $EF \parallel QR$  [By the converse of Basic Proportionality Theorem]

3. Let  $AB$  be a vertical pole of length 6m and  $BC$  be its shadow and  $DE$  be tower and  $EF$  be its shadow. Join  $AC$  and  $DF$ .

Now, in  $\triangle ABC$  and  $\triangle DEF$ , we have

$$\begin{aligned} \angle B &= \angle E = 90^\circ \\ \angle C &= \angle F && \text{(Angle of elevation of the Sun)} \\ \therefore \triangle ABC &\sim \triangle DEF && \text{(By AA criterion of similarity)} \end{aligned}$$

$$\text{Thus, } \frac{AB}{DE} = \frac{BC}{EF}$$

$$\Rightarrow \frac{6}{h} = \frac{4}{28} \quad (\text{Let } DE = h)$$

$$\Rightarrow \frac{6}{h} = \frac{1}{7} \quad \Rightarrow h = 42$$

$h = 42$  Hence, height of tower,  $DE = 42$ m




Fig. 7.12

4.








Fig. 7.13

Firstly, in  $\triangle ABC$ , we have

$LM \parallel CB$  (Given)

Therefore, by Basic Proportionality Theorem, we have

$$\frac{AM}{AB} = \frac{AL}{AC} \quad \dots(i)$$

Again, in  $\triangle ACD$ , we have

$LN \parallel CD$  (Given)

$\therefore$  By Basic Proportionality Theorem, we have

$$\frac{AN}{AD} = \frac{AL}{AC} \quad \dots(ii)$$

Now, from (i) and (ii), we have  $\frac{AM}{AB} = \frac{AN}{AD}$ .

5. In  $\triangle POQ$ , we have

$DE \parallel OQ$  (Given)

$\therefore$  By Basic Proportionality Theorem, we have

$$\frac{PE}{EQ} = \frac{PD}{DO} \quad \dots(i)$$

Similarly, in  $\triangle POR$ , we have

$DF \parallel OR$  (Given)

$$\frac{PD}{DO} = \frac{PF}{FR} \quad \dots(ii)$$

Now, from (i) and (ii), we have

$$\frac{PE}{EQ} = \frac{PF}{FR} \Rightarrow EF \parallel QR$$





Fig. 7.14

[Applying the converse of Basic Proportionality Theorem in  $\triangle PQR$ ]

6.



Given:  $\triangle ABC$  in which D and E are the mid-points of sides AB and AC respectively.

To prove:  $DE \parallel BC$

Proof: Since D and E are the mid-points of AB and AC respectively

$\therefore AD = DB$  and  $AE = EC$

$$\Rightarrow \frac{AD}{DB} = 1 \quad \text{and} \quad \frac{AE}{EC} = 1$$

$$\Rightarrow \frac{AD}{DB} = \frac{AE}{EC}$$

$DE \parallel BC$  Therefore,  $DE \parallel BC$  (By the converse of Basic Proportionality Theorem)

7. (i) In  $\triangle ABC$  and  $\triangle QRP$ , we have

$$\frac{BC}{RP} = \frac{2.5}{5} = \frac{25}{50} = \frac{1}{2}$$

$$\text{Hence, } \frac{AB}{QR} = \frac{AC}{QP} = \frac{BC}{RP}$$

$\therefore \triangle ABC \sim \triangle QRP$ , by SSS criterion of similarity.

(ii) In  $\triangle LMP$  and  $\triangle FED$ , we have

$$\frac{LP}{FD} = \frac{3}{6} = \frac{1}{2}, \quad \frac{MP}{ED} = \frac{2}{4} = \frac{1}{2}, \quad \frac{LM}{FE} = \frac{2.7}{5}$$

$$\text{Hence, } \frac{LP}{FD} = \frac{MP}{ED} \neq \frac{LM}{FE}$$

$\therefore \triangle LMP$  is not similar to  $\triangle FED$ .

(iii) In  $\triangle NML$  and  $\triangle PQR$ , we have

$$\angle M = \angle Q = 70^\circ$$

$$\text{Now, } \frac{MN}{QP} = \frac{2.5}{6} = \frac{5}{12} \quad \text{and} \quad \frac{ML}{QR} = \frac{5}{10} = \frac{1}{2}$$

$$\text{Hence, } \frac{MN}{QP} \neq \frac{ML}{QR}$$

$\triangle NML$  is not similar to  $\triangle PQR$ .

8.

In  $\triangle AOB$  and  $\triangle COD$ , we have

$$\angle AOB = \angle COD \quad [\text{Vertically opposite angles}]$$

$$\Rightarrow \frac{AO}{OC} = \frac{BO}{OD} \quad [\text{Given}]$$

So, by SAS criterion of similarity, we have

$$\triangle AOB \sim \triangle COD$$

$$\Rightarrow \frac{AO}{OC} = \frac{BO}{OD} = \frac{AB}{DC} \quad \Rightarrow \quad \frac{1}{2} = \frac{5}{DC} \quad [\because AB = 5 \text{ cm}]$$

$$\Rightarrow DC = 10 \text{ cm.}$$





Fig. 7.17

9.




In  $\triangle ABE$  and  $\triangle CFB$ , we have

$$\angle AEB = \angle CBF \quad (\text{Alternate angles})$$

$$\angle A = \angle C \quad (\text{Opposite angles of a parallelogram})$$

$$\therefore \triangle ABE \sim \triangle CFB \quad (\text{By AA criterion of similarity})$$

10.



In  $\triangle RPQ$  and  $\triangle RTS$ , we have


$$\angle RPQ = \angle RTS \text{ (Given)}$$

$$\angle PRQ = \angle TRS = \angle R \text{ (Common)}$$

$$\therefore \triangle RPQ \sim \triangle RTS \text{ (By AA criterion of similarity)}$$

### Long Answer :

1.



Given: A  $\triangle ABC$  in which D is the mid-point of AB and DE is drawn parallel to BC, which meets AC at E.

To prove:  $AE = EC$

Proof: In  $\triangle ABC$ ,  $DE \parallel BC$

$\therefore$  By Basic Proportionality Theorem, we have

$$\frac{AD}{DB} = \frac{AE}{EC} \dots (i)$$

Now, since D is the mid-point of AB

$$\Rightarrow AD = DB \dots (ii)$$

From (i) and (ii), we have

$$\frac{AD}{DB} = \frac{AE}{EC}$$





$$\Rightarrow 1 = \frac{AAND}{ANDC}$$

Hence, E is the mid-point of AC.

2. Given: ABCD is a trapezium, in which  $AB \parallel DC$  and its diagonals intersect each other at point O.

**To prove:**  $\frac{AO}{BO} = \frac{CO}{DO}$

**Construction:** Through O, draw  $OE \parallel AB$  i.e.,  $OE \parallel DC$ .

**Proof:** In  $\triangle ADC$ , we have  $OE \parallel DC$  (Construction)

$\therefore$  By Basic Proportionality Theorem, we have

$$\frac{AE}{ED} = \frac{AO}{CO} \quad \dots(i)$$

Now, in  $\triangle ABD$ , we have  $OE \parallel AB$  (Construction)

$\therefore$  By Basic Proportionality Theorem, we have

$$\frac{ED}{AE} = \frac{DO}{BO} \Rightarrow \frac{AE}{ED} = \frac{BO}{DO} \quad \dots(ii)$$

From (i) and (ii), we have

$$\frac{AO}{CO} = \frac{BO}{DO} \Rightarrow \frac{AO}{BO} = \frac{CO}{DO}$$




Fig. 7.35

3. In  $\triangle ABD$  and  $\triangle PQM$  we have

$$\angle B = \angle Q (\because \triangle ABC \sim \triangle PQR) \dots (i)$$

$$\frac{AB}{PQ} = \frac{BC}{QR} \quad (\because \triangle ABC \sim \triangle PQR)$$

$$\Rightarrow \frac{AB}{PQ} = \frac{\frac{1}{2} BC}{\frac{1}{2} QR} \Rightarrow \frac{AB}{PQ} = \frac{BD}{QM} \quad \dots(ii)$$




Fig. 7.36

[Since AD and PM are the medians of  $\triangle ABC$  and  $\triangle PQR$  respectively]

From (i) and (ii), it is proved that

$$\triangle ABD \sim \triangle PQM$$

(By SAS criterion of similarity)

$$\Rightarrow \frac{AB}{PQ} = \frac{BD}{QM} = \frac{AD}{PM} \Rightarrow \frac{AB}{PQ} = \frac{AD}{PM}$$

4. In  $\triangle EDC$  and  $\triangle EBA$  we have

$$\angle 1 = \angle 2 \text{ [Alternate angles]}$$

$$\angle 3 = \angle 4 \text{ [Alternate angles]}$$



$\angle CED = \angle AEB$  [Vertically opposite angles]

$\therefore \triangle EDC \sim \triangle EBA$  [By AA criterion of similarity]

$$\Rightarrow \frac{ED}{EB} = \frac{EC}{EA} \Rightarrow \frac{ED}{EC} = \frac{EB}{EA} \quad \dots(i)$$

It is given that  $\triangle AED \sim \triangle BEC$

$$\therefore \frac{ED}{EC} = \frac{EA}{EB} = \frac{AD}{BC} \quad \dots(ii)$$

From (i) and (ii), we get

$$\frac{EB}{EA} = \frac{EA}{EB} \Rightarrow (EB)^2 = (EA)^2 \Rightarrow EB = EA$$

Substituting  $EB = EA$  in (ii), we get

$$\frac{EA}{EA} = \frac{AD}{BC} \Rightarrow \frac{AD}{BC} = 1 \Rightarrow AD = BC$$





Fig. 7.37

5.



Given: A  $\triangle ABC$  in which  $\angle ABC = 90^\circ$  and  $AB = BC$ .

$\triangle ABD$  and  $\triangle CAE$  are equilateral triangles.

To Prove:  $\text{ar}(\triangle ABD) = \frac{1}{2} \times \text{ar}(\triangle CAE)$

Proof: Let  $AB = BC = x$  units.

$\therefore$  hyp.  $CA = \sqrt{x^2 + x^2} = x\sqrt{2}$  units.

Each of the  $\triangle ABD$  and  $\triangle CAE$  being equilateral has each angle equal to  $60^\circ$ .


$\therefore \triangle ABD \sim \triangle CAE$

But, the ratio of the areas of two similar triangles is equal to the ratio of the squares of their corresponding sides.

$$\therefore \frac{\text{ar}(\triangle ABD)}{\text{ar}(\triangle CAE)} = \frac{AB^2}{CA^2} = \frac{x^2}{(x\sqrt{2})^2} = \frac{x^2}{2x^2} = \frac{1}{2}$$

$$\text{Hence, } \text{ar}(\triangle ABD) = \frac{1}{2} \times \text{ar}(\triangle CAE)$$

6.



Given: Two triangles ABC and DEF, such that

$\triangle ABC \sim \triangle DEF$  and  $\text{area}(\triangle ABC) = \text{area}(\triangle DEF)$

To prove:  $\triangle ABC \cong \triangle DEF$

Proof:  $\triangle ABC \sim \triangle DEF$

$$\Rightarrow \angle A = \angle D, \angle B = \angle E, \angle C = \angle F$$

and 
$$\frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF}$$

Now, 
$$\text{ar}(\triangle ABC) = \text{ar}(\triangle DEF) \quad (\text{Given})$$

$$\therefore \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle DEF)} = 1$$

and 
$$\frac{AB^2}{DE^2} = \frac{BC^2}{EF^2} = \frac{AC^2}{DF^2} = \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle DEF)} \quad (\because \triangle ABC \sim \triangle DEF)$$

From (i) and (ii), we have

$$\frac{AB^2}{DE^2} = \frac{BC^2}{EF^2} = \frac{AC^2}{DF^2} = 1 \Rightarrow \frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF} = 1$$

$$AB = DE, BC = EF, AC = DF$$

$\triangle ABC \cong \triangle DEF$  (By SSS criterion of congruency)

7. Let  $\triangle ABC$  and  $\triangle PQR$  be two similar triangles. AD and PM are the medians of  $\triangle ABC$  and  $\triangle PQR$  respectively.

**To prove:**  $\frac{ar(\triangle ABC)}{ar(\triangle PQR)} = \frac{AD^2}{PM^2}$

**Proof:** Since  $\triangle ABC \sim \triangle PQR$

$$\frac{ar(\triangle ABC)}{ar(\triangle PQR)} = \frac{AB^2}{PQ^2}$$

...(i)




Fig. 7.40

In  $\triangle ABD$  and  $\triangle PQM$

$$\frac{AB}{PQ} = \frac{BD}{QM}$$

$$\left( \because \frac{AB}{PQ} = \frac{BC}{QR} = \frac{\frac{1}{2}BC}{\frac{1}{2}QR} \right)$$

and  $\angle B = \angle Q$

( $\because \triangle ABC \sim \triangle PQR$ )

Hence,  $\triangle ABD \sim \triangle PQM$

(By SAS similarity criterion)

$$\frac{AB}{PQ} = \frac{AD}{PM}$$

...(ii)

From (i) and (ii), we have

$$\frac{ar(\triangle ABC)}{ar(\triangle PQR)} = \frac{AD^2}{PM^2}$$

8.




Fig. 7.41

Join OA, OB and OC.

(i) In right  $\triangle$ 's OFA, ODB and OEC, we have

$$OA^2 = AF^2 + OF^2 \dots (i)$$

$$OB^2 = BD^2 + OF^2 \dots (ii)$$

$$OC^2 = EC^2 + OE^2$$

Adding (i), (ii) and (iii), we have

$$\Rightarrow OA^2 + OB^2 + OC^2 = AF^2 + BD^2 + EC^2 + OF^2 + OF^2 + OE^2$$

$$\Rightarrow OA^2 + OB^2 + OC^2 - OF^2 - OE^2 - OF^2 = AF^2 + BD^2 + EC^2$$






(ii) We have,  $OA^2 + OB^2 + OC^2 - IP^2 - OE^2 - OF^2 = AF^2 + BD^2 + EC^2$

$$\Rightarrow (OA^2 - OE^2) + (OB^2 - OF^2) - (OC^2 - IP^2) = AF^2 + BD^2 + EC^2$$

$$\Rightarrow AE^2 + CD^2 + BF^2 = AP^2 + BD^2 + EC^2$$

[Using Pythagoras Theorem in  $\triangle AOE$ ,  $\triangle BOF$  and  $\triangle COD$ ]

9.



**Fig. 7.42**

We have,  $DB = 3CD$

Now,

$$BC = BD + CD$$

$$\Rightarrow BC = 3CD + CD = 4CD \text{ (Given } DB = 3CD\text{)}$$

$$\therefore CD = \frac{1}{4} BC$$

$$\text{and } DB = 3CD = \frac{3}{4} BC$$

Now, in right-angled triangle ABD using Pythagoras Theorem we have

$$AB^2 = AD^2 + DB^2 \dots (i)$$

Again, in right-angled triangle  $\triangle ADC$ , we have

$$AC^2 = AD^2 + CD^2 \dots (ii)$$

Subtracting (ii) from (i), we have

$$AB^2 - AC^2 = DB^2 - CD^2$$


$$\Rightarrow AB^2 - AC^2 = \left(\frac{3}{4} BC\right)^2 - \left(\frac{1}{4} BC\right)^2 = \left(\frac{9}{16} - \frac{1}{16}\right) BC^2 = \frac{8}{16} BC^2$$

$$\Rightarrow AB^2 - AC^2 = \frac{1}{2} BC^2$$

$$\therefore 2AB^2 - 2AM^2 = BC^2$$

$$\Rightarrow 2AB^2 = 2AM^2 + BC^2$$

10.



Let ABC be an equilateral triangle and let  $AD \perp BC$ .

$$\therefore BD = DC$$

Now, in right-angled triangle ADB, we have

$$AB^2 = AD^2 + BD^2 \text{ [Using Pythagoras Theorem]}$$

$$\Rightarrow AB^2 = AD^2 + \left(\frac{1}{2} BC\right)^2 \Rightarrow AB^2 = AD^2 + \frac{1}{4} BC^2$$

$$\Rightarrow AB^2 = AD^2 + \frac{AB^2}{4} \quad [\because AB = BC]$$

$$\Rightarrow AB^2 - \frac{AB^2}{4} = AD^2 \Rightarrow \frac{3AB^2}{4} = AD^2 \Rightarrow 3AB^2 = 4AD^2$$

### Case Study Answers:

1. Answer :

i	c	$90^\circ$
ii	b	SAS
iii	b	4 : 9
iv	d	The converse of Pythagoras theorem
v	a	$48 \text{ cm}^2$

2. Answer :